

**ORAL PRESENTATION****Open Access**

# Strain-encoded cardiac magnetic resonance during high-dose dobutamine stress testing for the estimation of cardiac outcomes. Comparison to clinical parameters and conventional wall motion readings

Gitsios Gitsioudis Sr\*, Stephanie Lehrke, Nina Riedle, Sebastian J Buss, Evangelos Giannitsis, Nael F Osman, Hugo A Katus, Grigorios Korosoglou

From 2011 SCMR/Euro CMR Joint Scientific Sessions  
Nice, France. 3-6 February 2011

## Introduction

DS-MRI is a well established modality for the diagnostic classification and the risk stratification of patients with coronary artery disease (CAD). However, currently the detection of inducible ischemia by DS-MRI is based on the assessment of cine images, which is subjective and depends on the experience of the readers. In this regard, the human eye primarily tracks the radial displacement of the myocardium with cine-images, which is less sensitive than circumferential and longitudinal components for the detection of myocardial dysfunction.

## Purpose

To determine the prognostic value of Strain-Encoded-MRI (SENC) during high-dose dobutamine stress cardiac magnetic resonance imaging (DS-MRI) compared with conventional wall motion analysis.

## Methods

320 consecutive patients with suspected or known CAD underwent DS-MRI, using a standard protocol in a 1.5T MR-scanner. Wall motion and SENC were assessed at baseline and during stress, and outcome data including cardiac deaths, nonfatal myocardial infarctions ('hard events') and revascularization procedures performed >90 days after the MR-scans were collected.

## Results

35 hard events occurred during a 28±9 month follow-up period, including 10 cardiac deaths and 25 nonfatal myocardial infarctions, while 32 patients underwent coronary revascularization. Using multivariable Cox proportional-hazards models, atherogenic risk factors and inducible wall motion abnormalities (WMA) were independently associated with hard events (HR=1.5 (95% CI=1.1-2.1), p=0.02 for the total number risk factors and HR=3.9 (95%CI=1.8-8.6) for inducible WMA, p<0.001, respectively). When inducible strain defects were additionally considered in the model, however inducible WMA were no longer predictive, and SENC yielded the strongest independent association with cardiac outcome (HR of 10.5 (95%CI=2.9-38.5), p<0.001) (Figure 1). Furthermore, quantification analysis showed that a cut-off value of <1.64 for strain rate reserve was highly predictive for future hard cardiac events (AUC=0.82, 95% CI=0.78-0.86; negative predictive value=98%).

## Conclusions


Strain-Encoded-MRI aids the accurate identification of patients at high risk for future cardiac events and revascularization procedures, beyond the assessment of conventional atherogenic risk factors and inducible WMA on cine-images.

Published: 2 February 2011

University Clinic of Heidelberg, Heidelberg, Germany



© 2011 Gitsioudis et al; licensee BioMed Central Ltd. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Figure 1** Multivariable value of clinical parameters, baseline and stress magnetic resonance.

doi:10.1186/1532-429X-13-S1-O6

**Cite this article as:** Gitsioudis *et al.*: Strain-encoded cardiac magnetic resonance during high-dose dobutamine stress testing for the estimation of cardiac outcomes. Comparison to clinical parameters and conventional wall motion readings. *Journal of Cardiovascular Magnetic Resonance* 2011 **13**(Suppl 1):O6.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

